

POULTRY Form  
Management

SVC & GOVT degree college

\* Planning of poultry :-

\* Best practice management for meat chicken production in NSW

\* Better SIFR stewardship for meat poultry developments

\* poultry keeping environment :-

Information on the internal environment of sheds in which poultry are kept - brooder lighting fans feeders drinkers insulation etc

Information also on the external environment - the poultry farm and surrounding areas - noise duststock odour, development applications

\* Egg production systems in Australia

\* Infra-red brooding of intensive livestock

\* Alternative litter materials for poultry

\* Odour management options for meat chicken farms

\* Guard animals for livestock protection existing and potential use in Australia

## \* poultry management :-

\* Poultry management usually refers to the husbandry practices or production techniques that help to maximize the efficiency of production. Sound management practices in poultry are very essential to optimize production. Scientific poultry management aims at maximizing returns with minimum investment.

## \* poultry farming :-

\* Poultry farming is a form such as chickens, ducks, turkeys and geese raises domesticated birds to produce meat or eggs. Poultry - mostly chickens are farmed in great numbers.

In the United States, the national organization overseeing Poultry Production is the Food and Drug Administration (FDA). In UK, the national organization is the Department for Environment, Food and Rural Affairs (Defra).

\* Poultry farm in South Africa showing bare terrain new growth of nutritious grass

\* Chick :-

— — \* plural noun chicks :-

1. a young bird especially one newly hatched  
the egreher and reared two chicks

\* In formal :-

a young woman she is a great-looking chick  
Find here details of companies selling poultry  
for chicks for your purchase requirements Get  
lates in for no poultry farm chicks Country  
chicks will walk in it tracking litter and droppings  
that spread disease They tend to get wet  
and chilled and the stress will open the  
way to disease Some chicks may die  
by spilled water or leaky  
water are to be avoided.

## \* Broiler :-

A broiler is any chicken (*Gallus gallus domesticus*) that is bred and raised specifically for meat production most commercial broilers

reach slaughter weight between four and seven weeks of age although slower growing breeds reach slaughter

weight at approximately 14 weeks of age

\* Typical broilers have white feathers and yellowish skin Broiler or sometimes broil is also used sometimes to refer to younger chickens

under 2.0 kilograms (4.4 lb) as

compared with the larger roasters

## \* Broiler

Distribution world-wide

use meat

Traits

Skin color yellow

Classification Poultry

## \* Laying chickens :-

Laying chicken poultry farming means raising egg

laying Poultry bird for the purpose of commercial

egg production layer chickens are such a special species of hens, which need not be old. They remain laying eggs continuously till their 72-78 weeks of age.

They can produce about one kg of egg by

consuming about 2.25 kg of food during their egg laying period for the highly egg productive breeds available throughout the world.

## \* White egg laying Hens :-

This type of hens are comparatively smaller in size. Relatively eat less food and the color of egg shell is white. Cuckoo - BV - 300, Harvard, White Bovanish with etc are some popular chickens.

## \* Broiler medicine :-

\* Respiratory Herbs

\* Powerful Herbal Remedy for

[CRD - E-call]

\* Benefits for Birds / Animals

\* For prevention and treatment of chronic

Disease (CRD) and associated symptoms

including nasal discharge tracheal Cough Inflammation etc

\* Highly effective for treatment of

Mycoplasma Colibaculum Sethe call for chikawa Drink  
when Past mate

Poultry (CRD) e coli medicine Packaging type: ktr

500ml Packaging size: 1ltr 500ml

## \* Product Description :-

\* Packaging Type: (ktr 500ml)

Packaging Size: 1ltr 500ml

Dose/strength :- 5ml

Usage :- Clinical, Hospital

## \* Poultry Pump feeding :-

A starting point for how much to feed chickens each day. However, there is a simple figure to

provide you with a solid starting point  $\frac{1}{4}$  of a pound per fully grown chicken per day.

This means each chicken will approximate 1.5 pounds of feed in a week. Many of our feeds are sold in 40 lb

## \* Broilers :-

### \* Caring Broilers :-

Broilers are young chicken of either sex 10 weeks of age. Provide a mat with soft pliable smooth skin and breast bone cartilage.

### \* Housing :-

Provide 920 young chicken of either sex of six to eight weeks of age. Provide a mat with good ventilation. The general management of broiler is similar to those under egg type chicks.



## \* Feeding :-

Provide up to 2 weeks 35 cm and from  
Rais if tube feeders are used provide nos  
of 12kg Capacity feeders per 100 chicks

## \* Composition of broiler ration:-

Ingredient	Percentage Inclusion	
	Stare 0-5 weeks	Finger (6-7 weeks)
Yellow maize	47.00	54.50
Rice Polish	8.00	10.00
Soya bean meal	17.50	14.00
Groundnut cake (exp)	15.00	11.00
un salt dried fish	10.00	8.00
Mineral mixture	2.00	2.00
Salt	0.50	0.50
	100.00	100.00

Alternatively commercial broiler started and finisher rations prepared by reputed feed manufacturers can be given.

Names

S. Majunath Reddy

Sai Kumar

Nikil

C. Siva

Pranai Tej

M. Vara Prasad